

The Fragmentation Function of Gluon Splitting into P-wave Spin-singlet Heavy Quarkonium

Gang Hao¹, YaBing Zuo^{1,2}, Cong-Feng Qiao^{*1,3}

¹*Department of Physics, Graduate University, the Chinese Academy of Sciences
YuQuan Road 19A, Beijing 100049, China*

²*Department of Physics, Liaoning Normal University, Dalian 116029, China*

³*Theoretical Physics Center for Science Facilities (TPCSF), CAS,
YuQuan Road 19B, Beijing 100049, China*

Abstract

The gluon fragmentation function of $g \rightarrow h_c$ at leading order of strong coupling constant α_s and typical velocity v is calculated in the framework of NRQCD, of which the contributions from both color-singlet and -octet processes are taken into account and hence the result is infrared safe. It is found that the fragmentation probability of a high energy virtual gluon to split into h_c is about 7.1×10^{-7} . The analytic fragmentation function presented in this work can be employed to the study of h_c production in large transverse momentum at hadron colliders, such as Large Hadron Collider(LHC), and is also applicable for the study of h_b hadronproduction.

PACS number(s): 13.85.Ni, 12.38.Bx, 14.40.Pq

* qiaocf@gucas.ac.cn, corresponding author

It is well-known that the fragmentation mechanism dominates the heavy quarkonium hadroproduction at large transverse momentum [1, 2]. It is hence important to obtain the corresponding fragmentation function in order to properly estimate the production rate of a specific charmonium state. Fortunately it was found that these fragmentation functions for heavy quarkonium production are analytically calculable by virtue of perturbative quantum chromodynamics(QCD), with limited universal(phenomenological) parameters.

In past years, Braaten *et al.* calculated the fragmentation functions of gluon to the S-wave charmonium states η_c and J/ψ [1, 3]. Later, the gluon fragmentation to polarized J/ψ was obtained by fitting [4], and the fragmentation function of gluon to J/ψ via color-octet mechanism was calculated to next-to-leading(NLO) order accuracy [5]. The fragmentation functions of gluon to P-wave spin-triplet charmonium states χ_{cJ} were found infrared cut dependent [6], and after including the color-octet mechanism they then become infrared safe [2]. The polarized fragmentation functions of gluon to polarized χ_{cJ} were also obtained [7]. The fragmentation functions for gluon to D-wave spin-singlet and -triplet were presented in [8, 9]. The fragmentation functions of heavy quark splitting into S-wave charmonium states were calculated in [10, 11, 12], and the corresponding calculations for higher excited states, the P-wave and D-wave states, were given in Refs.[13, 14, 15, 16]. Notice that since in this letter we are not going to give a complete review of the calculation of quarkonium fragmentation functions, readers who are interested in the details on this respect should refer to recent review articles, like [17].

From above brief introduction, one may notice that the fragmentation function of gluon splitting into the spin-singlet P-wave state h_c is still absent. Because the production rate for h_c is low and the decay modes of h_c are obscure, which hinder experimenters to measure it, for a long time the study of h_c hadroproduction in large transverse momentum is not so urgent and hence the fragmentation function for it. However, in recently the h_c is observed in experiment [18, 19, 20, 21], and next people may further study its detailed natures. With the development of relevant experiments, people expect that more data on h_c will be collected, like in hadron collision, the LHC for example, in the near future [22, 23, 24]. In this sense, the fragmentation function of gluon splitting into h_c is necessary and valuable. In this paper we calculate it at the leading order of α_s and v in the framework of non-relativistic QCD(NRQCD) [25, 26], including contributions from both color-singlet and color-octet.

Generally speaking, heavy quarkonium hadroproduction via fragmentation mechanism involves both short-distance and long-distance effects. The differential cross section can be expressed as a convoluted form [27],

$$d\sigma_H(p) \approx \sum_i \int_0^1 dz d\hat{\sigma}_i(p/z, \mu) D_{i \rightarrow H}(z, \mu) . \quad (1)$$

This is a factorized formula, in which the short- and long-distance effects have been separated. $d\hat{\sigma}$ is the cross section for parton i production, and is perturbative QCD(pQCD) calculable due to the energetic parton and large momentum transfer. $D_{i \rightarrow H}(z, \mu)$ is a process-independent fragmentation function, which describes how parton i with invariant mass μ splitting into hadron H with light-cone momentum fraction z . For light hadrons, like ϕ and π , the fragmentation processes involve low energy scales, such as Λ_{QCD} and $m_{\phi, \pi}$, the fragmentation function is therefore a non-perturbative quantity and cannot be computed within the scope of pQCD. However, for the fragmentation process of a parton splitting into heavy quarkonium, though there will be some non-perturbative effects due to hadronization, the production of a heavy quark is a short-distance process and hence is pQCD accessible. It turns out therefore that the fragmentation function for heavy quarkonium production can be further factorized [3].

In the framework of non-relativistic QCD, the fragmentation function of a virtual gluon splitting into heavy quarkonium H reads

$$D_{g \rightarrow H}(z, \mu) = \sum_n d_n(z, \mu) \langle 0 | \mathcal{O}_n^H | 0 \rangle . \quad (2)$$

Here, n is the color-spin-orbital quantum number of the heavy quark pair with null relative momentum. The short-distance coefficient $d_n(z, \mu)$, describing the production of heavy quark pair with appropriate quantum number n , is pQCD calculable. $\mathcal{O}_n^H$ are local four-fermion operators in NRQCD, and their vacuum expectation values are proportional to the probabilities of heavy quark pairs with quantum number n hadronizing into quarkonium states. According to NRQCD, in (2) the short distance sector $d_n(z, \mu)$ can be computed order-by-order in strong coupling $\alpha_s(2m_c)$, and the long distance sector, the matrix elements, can be expanded in series of the typical relative velocity v of heavy quarks inside heavy quarkonium [3, 26]. In principle the fragmentation function is calculable to any order in α_s and v as desired; in practice, normally the NLO results are enough for phenomenological aim. It should be mentioned that in recently there are discussions about whether the definition




FIG. 1: Feynmann diagrams of the color-singlet process $g^* \rightarrow c\bar{c}[{}^1P_1^{(1)}] + g + g$




FIG. 2: Feynmann diagrams of the color-octet process $g^* \rightarrow c\bar{c}[{}^1S_0^{(8)}] + g$

of fragmentation function beyond NLO in strong coupling expansion is complete or not in the framework of NRQCD [28, 29], whereas it has no influence on our study in this work.

For process of gluon splitting into h_c , there are two classes of processes at leading order of α_s and v . One is the color-singlet process as shown in Fig.1, in which $c\bar{c}$ pair is in spin-singlet, color-singlet and P-wave state (denoted by ${}^1P_1^{(1)}$); the other is color-octet process as shown in Fig.2, in which the quark pair is in spin-singlet, color-octet and S-wave state (${}^1S_0^{(8)}$). The full fragmentation function for $g \rightarrow h_c$ then composes of color-singlet and color-octet terms, like

$$D_{g \rightarrow h_c}(z, \mu) = d_1(z, \Lambda) \langle 0 | \mathcal{O}_1^{h_c}({}^1P_1) | 0 \rangle + d_8(z) \langle 0 | \mathcal{O}_8^{h_c}({}^1S_0) | 0 \rangle (\Lambda), \quad (3)$$

where the short-distance coefficients $d_1(z, \Lambda)$ and $d_8(z)$ can be obtained through the cal-

ulation of Feynmann diagrams Fig.1 and Fig.2, respectively. The factorization scale Λ is introduced to separate the effect at short distance of order $1/m_c$ from the one at long distance of the radius of heavy quarkonium $1/(m_c v)$. $\langle 0|\mathcal{O}_1^{h_c}(^1P_1)|0\rangle$ and $\langle 0|\mathcal{O}_8^{h_c}(^1S_0)|0\rangle$ are matrix elements of NRQCD operators, scaling as $m_c^5 v^5$ and $m_c^3 v^5$ according to velocity-scaling rules in NRQCD. Their dependence on Λ can be obtained by renormalization group equations. To leading order of $\alpha_s(\Lambda)$ they are [26],

$$\begin{aligned}\Lambda \frac{d}{d\Lambda} \langle 0|\mathcal{O}_1^{h_c}(^1P_1)|0\rangle &= 0 \\ \Lambda \frac{d}{d\Lambda} \langle 0|\mathcal{O}_8^{h_c}(^1S_0)|0\rangle &= \frac{4C_F\alpha_s(\Lambda)}{3N_c\pi m_c^2} \langle 0|\mathcal{O}_1^{h_c}(^1P_1)|0\rangle.\end{aligned}\quad (4)$$

The last equations can be used to sum up large logarithms of m_c/Λ when the factorization scale are chosen to be much smaller than m_c [2]. i.e.,

$$\langle 0|\mathcal{O}_8^{h_c}(^1S_0)|0\rangle(m_c) = \langle 0|\mathcal{O}_8^{h_c}(^1S_0)|0\rangle(\Lambda) + \frac{4C_F}{3N_c m_c^2 \beta_0} \ln \frac{\alpha_s(\Lambda)}{\alpha_s(m_c)} \langle 0|\mathcal{O}_1^{h_c}(^1P_1)|0\rangle, \quad (5)$$

with QCD beta function $\beta_0 = (11N_c - 2n_f)/6$. In following analytical results, we will show that $d_8(z)$ is Λ independent at leading order of α_s . Thus, the dependence on Λ in the factorization formula (3) comes from $\langle 0|\mathcal{O}_8^{h_c}(^1S_0)|0\rangle$ and $d_1(z, \Lambda)$, the same as in the fragmentation functions of $g \rightarrow \chi_{cJ}$.

Leaving the polarization of fragmenting gluon unsummed, we obtain the squared amplitudes of color-singlet processes,

$$\begin{aligned}\sum \mathcal{A}_\mu^* \mathcal{A}_\nu &= F_1 g_{\mu\nu} + F_2 p_\mu p_\nu + F_3 (p_\mu k_\nu + k_\mu p_\nu) + F_4 k_\mu k_\nu \\ &+ (\text{terms proportional to } q_\mu \text{ or } q_\nu) .\end{aligned}\quad (6)$$

Here, p , k and q are the momenta of the $c\bar{c}$ pair, final and initial gluons, and $s = q^2$ is the virtuality of fragmenting gluon. In the condition $q_0 \gg m_c$, which is required to make fragmentation mechanism a reasonable approximation, the fragmenting gluon is almost transversely polarized. And under this condition we can drop terms proportional to q_μ or q_ν , which are considered as gauge artifact and suppressed in an appropriate axial gauge [1]. The Lorentz indices in Eq.(6) imply that there are angle dependence in the squared amplitudes. After averaging the directions of momentum p and k , we get the following replacements,

$$p_\mu p_\nu \rightarrow -\frac{1}{2} |\vec{p}_\perp|^2 g_{\mu\nu}, \quad (7)$$

$$k_\mu k_\nu \rightarrow -\frac{1}{2}g_{\mu\nu}\frac{t^2}{y^2-r}\left[\frac{1}{2}|\vec{p}_\perp|^2(3\cos^2\theta_k-1)+s(y^2-r)(1-\cos^2\theta_k)\right], \quad (8)$$

$$k_\mu p_\nu + p_\mu k_\nu \rightarrow -g_{\mu\nu}|\vec{p}_\perp|^2\frac{t\cos\theta_k}{\sqrt{y^2-r}}, \quad (9)$$

with

$$\cos\theta_k = \frac{(1+r)/2-t-y+ty}{t\sqrt{y^2-r}}, \quad |\vec{p}_\perp|^2 = s(-r+2yz-z^2), \quad (10)$$

where the dimensionless variables are defined,

$$y = \frac{p \cdot q}{s}, \quad t = \frac{k \cdot q}{s}, \quad r = \frac{4m_c^2}{s}, \quad z = \frac{p^0 + p^3}{q^0 + q^3}. \quad (11)$$

Notice that the squared amplitude of Eq.(6) is now proportional to metric tensor $g_{\mu\nu}$, and the fragmenting gluon behaves like an on-shell one.

To avoid the presence of infrared divergence while gluons in final state become soft, we set a lower cutoff Λ on the energy of the final gluons, then the color-singlet coefficient $d_1(z)$ converges and depends on Λ . After integrating over the three-body phase space, the color-singlet part of fragmentation function in Eq.(3) is obtained,

$$\begin{aligned} d_1(z, \Lambda) = & \frac{5\alpha_s^3(\mu)}{20736\pi m_c^5} \int_0^z dr \int_{(r+z^2)/2z}^{(1+r)/2} dy \frac{1}{(1-y)^4(y^2-r)^2} \left\{ \frac{(1+r^2)-4yz+2z^2}{(1-r)^2} h(y, r) \right. \\ & + \frac{1}{(y-r)^5} \sum_{i=0}^2 z^i \left((r-y)f_i(y, r) + \frac{g_i(y, r)}{\sqrt{y^2-r}} \ln \frac{y-r+\sqrt{y^2-r}}{y-r-\sqrt{y^2-r}} \right) \Big\} \\ & + \frac{5\alpha_s^3(\mu)}{162\pi m_c^5} \left\{ -[-2z^2+3z+2(1-z)\ln(1-z)] \ln \frac{\Lambda}{m_c} + w(z) \right\}, \end{aligned} \quad (12)$$

with functions $w(z)$, $h(y, r)$, $f_i(y, r)$ and $g_i(y, r)$ read

$$\begin{aligned} w(z) = & 2(1-z)\ln^2(1-z) + [-2-3(-2+z)z+3(-1+z)\ln z]\ln(1-z) \\ & + \frac{1}{6}[2\pi^2(1-z)+3z(-7+8z)+3z(-3+2z)\ln z] - 2(1-z)Li_2(1-z) - (1-z)Li_2(z), \end{aligned} \quad (13)$$

$$\begin{aligned} h(y, r) = & (1+11r-5r^2+r^3)(1-20r+6r^2-4r^3+r^4) + (2-20r+606r^2-280r^3+94r^4-20r^5+2r^6)y \\ & + (4+468r-792r^2+232r^3-44r^4+4r^5)y^2 + (8-1120r+560r^2-96r^3+8r^4)y^3 \\ & + (-240+1072r-208r^2+16r^3)y^4 + (544-448r+32r^2)y^5 + (-448+64r)y^6 + 128y^7, \end{aligned} \quad (14)$$

$$\begin{aligned} f_0(y, r) = & r^3(36-15r+25r^2-57r^3+59r^4-97r^5+23r^6-7r^7+r^8) \\ & + (-164r^3+54r^4+80r^5-34r^6+344r^7-2r^8+12r^9-2r^{10})y \\ & + (-82r^2+274r^3-374r^4-46r^5-746r^6+26r^7-14r^8+2r^9)y^2 \\ & + (-4r+488r^2+88r^3+652r^4+824r^5-564r^6-108r^7)y^3 \\ & + (1+33r-1277r^2-637r^3-877r^4+1947r^5+553r^6+r^7)y^4 \\ & + (2-160r+1734r^2+488r^3-2794r^4-1032r^5+66r^6)y^5 \end{aligned}$$

$$\begin{aligned}
& +(-28 + 356r - 1144r^2 + 2232r^3 + 644r^4 - 396r^5)y^6 + (136 - 352r - 352r^2 + 400r^3 + 968r^4)y^7 \\
& +(-336 + 80r - 1136r^2 - 1264r^3)y^8 + (448 + 736r + 992r^2)y^9 + (-384 - 448r)y^{10} + 128y^{11}, \tag{15}
\end{aligned}$$

$$f_1(y, r) = -2yf_2(y, r), \tag{16}$$

$$\begin{aligned}
f_2(y, r) = & 2[r^2(6 + 8r - 47r^2 + 33r^3 - 24r^4 + 46r^5 - 7r^6 + r^7) \\
& +(-16r^2 + 96r^3 + 56r^4 + 2r^5 - 206r^6 - 10r^7 - 2r^8)y \\
& +(-16r - 148r^2 - 248r^3 - 74r^4 + 502r^5 + 110r^6 + 2r^7)y^2 \\
& +(124r + 470r^2 + 286r^3 - 614r^4 - 234r^5 + 16r^6)y^3 \\
& +(-7 - 285r - 592r^2 + 320r^3 + 151r^4 - 99r^5)y^4 + (34 + 336r + 204r^2 + 128r^3 + 242r^4)y^5 \\
& +(-84 - 220r - 340r^2 - 316r^3)y^6 + (112 + 232r + 248r^2)y^7 + (-96 - 112r)y^8 + 32y^9], \tag{17}
\end{aligned}$$

$$\begin{aligned}
g_0(y, r) = & 2[r^4(-9 + r - 8r^2 - 8r^3 + r^4 - r^5 - 8r^6) + (46r^4 + 10r^5 + 71r^6 + 20r^7 + 4r^8 + 66r^9 + 7r^{10})y \\
& +(28r^3 - 111r^4 - 94r^5 - 148r^6 - 34r^7 - 213r^8 - 20r^9)y^2 \\
& +(-148r^3 + 193r^4 + 146r^5 + 150r^6 + 356r^7 - 111r^8 - 26r^9)y^3 \\
& +(-28r^2 + 297r^3 - 232r^4 - 120r^5 - 330r^6 + 695r^7 + 182r^8)y^4 \\
& +(192r^2 - 166r^3 + 452r^4 + 152r^5 - 1604r^6 - 498r^7 + 16r^8)y^5 \\
& +(-608r^2 - 590r^3 - 662r^4 + 1930r^5 + 614r^6 - 124r^7)y^6 \\
& +(1160r^2 + 1152r^3 - 896r^4 - 112r^5 + 408r^6)y^7 \\
& +(8r - 1152r^2 - 216r^3 - 696r^4 - 744r^5)y^8 + (-104r + 360r^2 + 864r^3 + 816r^4)y^9 \\
& +(8 + 216r - 376r^2 - 520r^3)y^{10} + (-16 - 48r + 160r^2)y^{11}], \tag{18}
\end{aligned}$$

$$g_1(y, r) = -2yg_2(y, r), \tag{19}$$

$$\begin{aligned}
g_2(y, r) = & 2[r^3(-3 - 5r + 8r^2 + 3r^4 + 13r^5 + 16r^6) - (-10r^3 - 14r^4 + 11r^5 + 4r^6 + 84r^7 + 138r^8 + 11r^9)y \\
& -(-10r^2 + 11r^3 + 114r^4 + 60r^5 - 196r^6 - 495r^7 - 76r^8)y^2 \\
& -(16r^2 - 183r^3 - 492r^4 + 86r^5 + 940r^6 + 201r^7 - 8r^8)y^3 \\
& -(16r + 175r^2 + 796r^3 + 640r^4 - 914r^5 - 215r^6 + 62r^7)y^4 \\
& -(-96r - 688r^2 - 1172r^3 + 108r^4 - 76r^5 - 204r^6)y^5 \\
& -(218r + 922r^2 + 562r^3 + 502r^4 + 372r^5)y^6 + (200r + 472r^2 + 568r^3 + 408r^4)y^7 \\
& -(-4 + 36r + 268r^2 + 260r^3)y^8 - (8 - 8r - 80r^2)y^9]. \tag{20}
\end{aligned}$$

Compared to the color-singlet short-distance coefficient, the calculation for the color-octet one is much more straightforward and the result does not depend on Λ ,

$$d_8(z) = \frac{5\alpha_s^2(\mu)}{96m_c^3}[-2z^2 + 3z + 2(1 - z)\ln(1 - z)]. \tag{21}$$

Now we have the final expression for fragmentation function $D_{g \rightarrow h_c}(z, \mu)$ at the leading order in strong coupling constant. If the value for factorization scale is chosen to be much smaller than m_c , we should exploit the renormalization group equation (4), or its solution (5), to sum up the logarithm of Λ/m_c . In this work, the factorization scale is set to be m_c so that we can avoid the large logarithms [2]. The value of the scale μ in α_s are chosen to be $2m_c$,

which is the square root of the minimum value of the virtuality of the fragmenting gluon.

Integrating the fragmentation function $D_{g \rightarrow h_c}(z, 2m_c)$ over the momentum fraction z , we get the fragmentation probability,

$$\int_0^1 dz D_{g \rightarrow h_c}(z, 2m_c) = \frac{5\alpha_s^2(2m_c)}{96m_c^3} \left[\frac{(-54.5)\alpha_s(2m_c)}{216\pi m_c^2} \langle \mathcal{O}_1^{h_c} \rangle + (0.33) \langle \mathcal{O}_8^{h_c} \rangle(m_c) \right]. \quad (22)$$

To make the total probability positive, the lower bound for the magnitude of color-octet matrix element at factorization scale $\Lambda = m_c$ is set to be

$$\langle \mathcal{O}_8^{h_c} \rangle(m_c) > \frac{3\alpha_s(2m_c)}{4\pi m_c^2} \langle \mathcal{O}_1^{h_c} \rangle. \quad (23)$$

For numerical evaluation, the input parameters are taken as follows: $m_c = m_{h_c}/2 = 1.78\text{GeV}$, $\alpha_s(2m_c) = 0.26$, $\langle 0 | \mathcal{O}_1^{h_c}(^1P_1) | 0 \rangle = 0.32\text{GeV}^5$ [30], which enables $\langle 0 | \mathcal{O}_8^{h_c}(^1S_0) | 0 \rangle(m_c) > 6.3 \times 10^{-3}\text{GeV}^3$, $\langle 0 | \mathcal{O}_8^{h_c}(^1S_0) | 0 \rangle(m_c) = 9.8 \times 10^{-3}\text{GeV}^3$ [24, 31]. With above inputs the fragmentation probability in Eq.(22) is found to be about 7.1×10^{-7} . Compared with the probabilities of gluon fragmenting into χ_{c0} , χ_{c1} , and χ_{c2} , which are 0.4×10^{-4} , 1.8×10^{-4} and 2.4×10^{-4} [2], the value of gluon fragmentation to h_c is smaller by two to three orders, and it is even less than the color-singlet probability of process $g^* \rightarrow J/\psi gg$ by an order [3]. The z dependence of the fragmentation function at $\mu = 2m_c$ and $\Lambda = m_c$ is shown in Figure 3.

In conclusion, we have computed in this work the fragmentation function of gluon to P-wave spin singlet quarkonium h_c . The analytic expression is infrared safe while both color-singlet and -octet processes are taken into account. It is found that the fragmentation probability of a high energy virtual gluon splitting into h_c is about 7.1×10^{-7} . The obtained analytic fragmentation function can be employed to the study of h_c hadroproduction, especially in the large transverse momentum region. Finally, it is worthy to note that result in this work can be readily applied to the study of h_b physics.

Acknowledgments

This work was supported in part by the National Natural Science Foundation of China(NSFC) under the grants 10935012, 10928510, 10821063, 10775179 and by CAS Key




FIG. 3: The fragmentation function $D_{g \rightarrow h_c}(z, \mu)$ as a function of z for $\mu = 2m_c$ and $\Lambda = m_c$, the dotted line is the contribution from color-singlet process, the dash-dotted line is the contribution from color-octet process and the solid line is the total result.

Projects (KJCX2-yw-N29, H92A0200S2).

-
- [1] E. Braaten and T.C. Yuan, Phys. Rev. Lett. **71**, 1673 (1993).
 - [2] E. Braaten and T.C. Yuan, Phys. Rev. D**50**, 3176 (1994).
 - [3] E. Braaten and T.C. Yuan, Phys. Rev. D**52**, 6627 (1995).
 - [4] W. Qi, C.F. Qiao, and J.X. Wang, Phys. Rev. D**75**, 074012 (2007).
 - [5] Eric Braaten and Jungil Lee, Nucl. Phys. B**586**, 427 (2000).
 - [6] J.P. Ma, Nucl. Phys. B**447**, 405 (1995).
 - [7] P. Cho, M.B. Wise and S.P. Trivedi, Phys. Rev. D**51**, 2039 (1995).
 - [8] P. Cho and M.B. Wise, Phys. Rev. D**51**, 3352 (1995).

- [9] C.-F. Qiao, F. Yuan and K.T. Chao, Phys. Rev. D**55**, 5437 (1997).
- [10] E. Braaten, K. Cheung and T. C. Yuan, Phys. Rev. D**48**, 4230(1993); Phys. Rev. D**48**, R5049 (1993).
- [11] C. H. Chang and Y. Q. Chen, Phys. Lett. B**284**, 127(1992); Phys. Rev. D**46**, 3845 (1992).
- [12] A.F. Falk, M. Luke, M.J. Savage and M.B. Wise, Phys. Lett. B**312**, 486 (1993); Phys. Rev. D**49**, 555 (1994)
- [13] J.P. Ma, Phys. Rev. D**53**, 1185 (1996).
- [14] Y. Q. chen, Phys. Rev. D**48**, 5158 (1993); H. Pois and T. C. Yuan, U.C.-Davis preprint UCD-94-2 (April, 1994).
- [15] T. C Yuan, Phys. Rev. D**50**, 5664 (1994).
- [16] King-man Cheung and Tzu Chiang Yuan, Phys. Rev. D**53**, 3591 (1996).
- [17] D.M. Asner, *et al.* preprint: IHEP-PHYSICS-REPORT-BES-III-2008-001; arXiv:0809.1869.
- [18] E835 Collaboration, M. Andreotti, *et al.*, Phys. Rev. D**72**, 032001 (2005).
- [19] CLEO Collaboration, J.L. Rosner, *et al.*, Phys. Rev. Lett. **95**, 102003 (2005).
- [20] CLEO Collaboration, P. Rubin, *et al.*, Phys. Rev. D**72**, 092004 (2005).
- [21] CLEO Collaboration, S. Dobbs, *et al.*, Phys. Rev. Lett. **101**, 182003 (2008).
- [22] Cong-Feng Qiao and Changzheng Yuan, Phys. Rev. D**63**, 014007 (2001).
- [23] K. Sridhar, Phys. Lett. B**674**, 36 (2009).
- [24] Cong-Feng Qiao, De-Long Ren and Peng Sun, Phys. Lett. B**680**, 159 (2009).
- [25] W.E. Caswell and G.P. Lepage, Phys. Lett. B**167**, 437 (1986).
- [26] G.T. Bodwin, E. Braaten and G.P. Lepage, Phys. Rev. D**51**, 1125(1995), Erratum-ibid. D**55**, 5853 (1997).
- [27] G. Curci, W. Furmanski and R. Petronzio, Nucl. Phys. B**175**, 27 (1985); J.C. Collins and D.E. Soper, Nucl. Phys. B**194**, 445 (1982).
- [28] Gouranga C. Nayak, Jian-Wei Qiu, and George Sterman, Phys. Lett. B**613**, 45 (2005).
- [29] Gouranga C. Nayak, Jian-Wei Qiu, and George Sterman, Phys. Rev. D**72**, 114012 (2005).
- [30] M. Mangano and A. Petrelli, Phys. Lett. B**352**, 445 (1995).
- [31] P.L. Cho and A.K. Leibovich, Phys. Rev. D**53**, 150 (1996); Phys. Rev. D**53**, 53 (1996).
- [32] J.H. Kuhn, J. Kaplan and E.G.O. Safiani, Nucl. Phys. B**157**, 125 (1979).